




Who is eligible for BioNTech bivalent vaccine booster ?






Did not infected with COVID-19 before

Age group		1st dose	Interval	2nd dose	Interval	3rd dose	Interval	4th dose
 12 - 17 years old	General Public	N/A		N/A		N/A		N/A
	Immunocompromised person	N/A		N/A		N/A	90 days after 3rd dose	✓
 18 - 49 years old	General Public	N/A		N/A	90 days after 2nd dose	✓	180 days after 3rd dose	✓
	Immunocompromised person	N/A		N/A		N/A	90 days after 3rd dose	✓
 ≥50 years old	General Public	N/A		N/A	90 days after 2nd dose	✓	90 days after 3rd dose	✓
	Immunocompromised person	N/A		N/A		N/A	90 days after 3rd dose	✓

* Person should have previously received at least two doses of the same type of COVID-19 vaccine

COVID-19 Recovered Persons

Age group		1st dose	Interval	2nd dose	Interval	3rd dose	Interval	4th dose
 12 - 17 years old	General Public	N/A		N/A		N/A		N/A
	Immunocompromised person	N/A		N/A	90 days after recovery or after 2nd dose (whichever is later)	✓		N/A
 18 - 49 years old	General Public	N/A	90 days after recovery or after 1st dose (whichever is later)*	✓	180 days after recovery or after 2nd dose (whichever is later)	✓	180 days after recovery or after 3rd dose (whichever is later)	✓
	Immunocompromised person	N/A		N/A		N/A	90 days after recovery or after 3rd dose (whichever is later)	✓
 ≥50 years old	General Public	N/A	90 days after recovery or after 1st dose (whichever is later)	✓	90 days after recovery or after 2nd dose (whichever is later)^	✓	180 days after recovery or after 3rd dose (whichever is later)	✓
	Immunocompromised person	N/A		N/A		N/A	90 days after recovery or after 3rd dose (whichever is later)	✓

✓ Eligible for BioNTech bivalent vaccine booster

#Immunocompromised persons include the following groups:

(i) Cancer or hematological malignancy on active immunosuppressive treatment now or in the past 12 months; (ii) Recipients of solid organ transplant or stem cell transplant on immunosuppressive treatment; (iii) Severe primary immunodeficiency or on chronic dialysis; (iv) Advanced or untreated HIV disease (v) On active immunosuppressive drugs, or immunosuppressive chemotherapy/radiotherapy in past 6 months

* If recovered persons have received 1 dose of COVID-19 vaccine before infection, they may receive their next dose for personal reasons at a shorter interval of no less than 28 days after recovery.

^ If recovered persons have received 2 doses of COVID-19 vaccine before infection, they may receive their next dose for personal reasons at a shorter interval of no less than 28 days after recovery.

Online Booking

More Details



Gleneagles Hospital

HONG KONG
港怡醫院

